细胞外囊泡及其在疾病诊疗中的应用

黄琳^{1,2†}, 王殿冰^{2†}, 顾宁¹, 张先恩^{1,2*}

1. 东南大学生物科学与医学工程学院, 南京 210096;

2. 中国科学院生物物理研究所, 中国科学院生物大分子研究中心, 生物大分子国家实验室, 北京 100101

† 同等贡献

* 联系人, E-mail: zhangxe@ibp.ac.cn

2019-03-11 收稿, 2019-03-27 修回, 2019-03-29 接受, 2019-06-20 网络版发表

国家重点研发专项(2017YFA0205500)资助

摘要 细胞外囊泡(extracellular vesicles, EVs)是细胞分泌的一种脂质囊泡, 尺寸介于30~5000 nm. 通过不同的发生机理, EVs可以形成微囊泡体(microvesicles)、凋亡小体(apoptotic bodies)和外泌体(exosomes)等. 它们通过携带母细胞的不同脂质、蛋白质和核酸等活性成分来靶向附近或远端细胞, 在细胞的信息交流及生理病理过程中均具有重要作用. EVs在生物体系内广泛存在, 其生物学功能也越来越多地被认识, 或许正在成为一个“细胞外囊泡生物学”领域. 对EVs性质的认识也促进了相关应用研究, 涉及疾病诊断、治疗和药物运送等. 本文重点阐述了EVs的生物起源、生物组成、生物学特性、生物学功能、制备和表征手段, 并针对EVs在疾病诊疗中的应用所面临的问题展开讨论. 可以预期, 对EVs形成和调控机理的深入认识, 一方面有助于我们更好地理解EVs的生物学功能、异质性和功能多样性; 另一方面有可能基于这些知识来解决EVs开发应用中的瓶颈问题.

关键词 细胞外囊泡, 药物运送载体, 外泌体, 纳米材料, 肿瘤标志物

细胞外囊泡(extracellular vesicles, EVs)是几乎所有细胞都能分泌的一种脂质囊泡, 外层由双层脂质膜包裹, 内部包含各种蛋白质和核酸, 尺寸30~5000 nm^[1]. EVs主要包括微囊泡体(50~1000 nm)、凋亡小体(50~5000 nm)和外泌体(30~150 nm), 分布于各种体液中(如血液、淋巴液、唾液、尿液以及乳汁等)^[2]. 1967年, 由磷脂组成的囊泡首次在血小板中被发现, 并被认为是细胞释放的一种垃圾^[3]. 1985年, Johnstone等人^[4]通过电子显微镜首次观察到网织红细胞中囊泡的结构. 随后, 不同类型的囊泡在各种细胞中相继被发现, 其生物学作用逐渐受到关注. 据检索Web of Science数据(Clarivate Analytics), 近10年来, 文献量迅速增长, 每年达数千篇, 涉及多个研究方向, 包括形成机制、组成成分、生理功能、应用潜力等. 本文将结合我们自己的

研究体会作一概述.

1 EVs的生物学特性

1.1 EVs的生物起源及其异质性

研究表明, 细胞可以分泌多种不同类型的EVs, 且每类EVs都具独特的生物发生机制, 存在特征性蛋白质和核酸^[5]. 根据生物起源, EVs可分为3类(图1): 微囊泡体(microvesicles)、凋亡小体(apoptotic bodies)以及外泌体(exosomes)^[2].

微囊泡体是由细胞质膜直接出芽形成, 尺寸介于50~1000 nm. 其释放受许多因素影响, 如细胞质Ca²⁺离子水平的变化^[6]、ADP核糖基化因子6 (ADP ribosylation factor 6, ARF6)的抑制^[7]、细胞质膜成分的变化^[8]

引用格式: 黄琳, 王殿冰, 顾宁, 等. 细胞外囊泡及其在疾病诊疗中的应用. 科学通报, 2019, 64: 2025~2036

Huang L, Wang D B, Gu N, et al. Extracellular vesicles and their application in the diagnosis and treatment of diseases (in Chinese). Chin Sci Bull, 2019, 64: 2025~2036, doi: 10.1360/N972019-00018

图1 (网络版彩色)细胞释放的几种主要EVs, 改编自文献[2]。其中, 微囊泡体、肿瘤小体和凋亡小体是由细胞质膜直接出芽形成, 外泌体是由多囊泡体(Multivesicular bodies, MVBs)与细胞质膜融合释放形成

Figure 1 (Color online) The main types of extracellular vesicles released by cells, adapted from ref. [2]. Microvesicles, oncosomes and apoptotic bodies were formed by direct budding of the plasma membrane, and exosomes were formed by the fusion of MVBs with the plasma membrane

和细胞骨架重组^[9]等。

微囊泡体还包括一种由肿瘤细胞释放的肿瘤小体(Oncosomes), 尺寸为1~10 μm, 由Janus Rak's研究组于2008年首次在胶质瘤细胞中发现^[10]。与其他EVs相比, 虽然肿瘤小体中CD63等蛋白的表达量较少, 但其可特异性富集细胞角蛋白-18(cytokeratin-18)等标志物, 此外还存在许多与肿瘤相关的蛋白质和核酸^[11]。肿瘤小体的释放也可通过多种方式进行诱导, 如过表达癌症相关蛋白(如Akt1, caveolin-1等)或活化特定代谢途径等^[2,12]。

凋亡小体是细胞遭受程序性死亡或凋亡晚期时释放的一种囊泡, 尺寸为50~2000 nm。凋亡小体主要通过随机包裹的方式把蛋白质等组分装入内腔中, 也有报道RNA等可以特异性地被分选到凋亡小体中^[2]。

外泌体是目前研究得最多的囊泡, 由细胞通过MVBs与细胞质膜融合释放到胞外, 尺寸介于30~150 nm。1987年, Johnstone等人^[13]首次提出了外泌体这个名词, 并给出定义。2013年, Rothman等3位科学家因发现了细胞囊泡运输调控机制获得诺贝尔生理学或医学奖, 外泌体的生物发生与囊泡运输调控密切相关, 引发广泛关注。以下, 我们将重点对其生物起源及其调控机制进行概述。

外泌体的形成过程主要有3个阶段, 细胞质膜内陷、MVBs形成以及MVBs与细胞质膜的融合释放^[14]。首先, 细胞质膜会通过内吞作用形成小泡, 继而与早期

内体(early endosomes)融合。然后, 早期内体发生内陷形成MVBs。在此过程中, 细胞质中的成分会通过不同的分选机理分选到管腔囊泡(intraluminal vesicles, ILVs)中。研究表明, ILVs主要通过内体分选复合物(endosomal sorting complex required for transport, ESCRT)依赖性和ESCRT非依赖性的机理进行货物分选^[2]。最后, MVBs与细胞质膜进行融合, 释放出外泌体。

ESCRT主要包括4种复合体(ESCRT-0, ESCRT-I, ESCRT-II, ESCRT-III)和一些辅助蛋白(VPS4, VTA1, ALIX等)^[15]。通过对23种ESCRT组分分析发现, 敲除一些蛋白(如HRS, STAM1, TSG101)可以降低外泌体的分泌^[16,17], 而有些组分(如VPS4)的作用却与之相反^[18]。又如, 辅助蛋白ALIX可通过与同线蛋白(syntenin)的相互作用促进外泌体的生成^[19]。可见, ESCRT家族成员对外泌体的生物起源的影响是复杂的。

ESCRT非依赖性调控机制与一些四跨膜蛋白(tetraspanins)、热休克蛋白(heat shock protein)和脂质等的调控高度相关^[15,20,21]。例如脂类调控机制中, 中性鞘磷脂酶(neutral sphingomyelinase)可以水解鞘磷脂生成神经酰胺, 而神经酰胺会影响MVBs形成^[22]。磷脂酶D2(phospholipase D2)水解生成的磷脂酸也可以像神经酰胺一样促进MVBs形成, 进而影响外泌体生成^[23]。而热休克蛋白HSP70主要是调控转铁蛋白受体(ferritin receptor)在外泌体上的富集^[24]。

MVBs形成后, 会与微丝和微管骨架相互作用转运至细胞质膜^[25,26]。其中, 皮动蛋白(cortactin)可以通过与微丝相关蛋白复合体Arp2/3以及微丝的相互作用来影响MVBs与细胞质膜的对接, 从而调控外泌体释放^[27]。研究表明, 一种小分子GTP结合蛋白, Ras超家族中的亚族RAB蛋白也参与此过程的调节^[28], 如Rab27a^[29,30], Rab27b^[31]和Rab35等^[32]。随后, 在MVBs与细胞质膜的融合过程中, 一些可溶性N-乙基马来酰亚胺敏感因子附着蛋白受体(soluble N-ethylmaleimide-sensitive factor attachment protein receptors, SNARE)会参与其调控过程^[33], 如重组人小泡相关膜蛋白VAMP7和重组人突触囊泡蛋白同源物YKT6。

1.2 EVs的组成及生物学功能

EVs的主要组成成分有蛋白质、脂质和核酸, 不同细胞来源和不同细胞状态下产生的EVs具有不同的内容物^[34]。在Vesiclepedia, EVpedia和ExoCarta这3个数据库中, 与EVs有关的成分已涉及上万种蛋白和RNA(包括

表 1 EVs的主要组成成分^[33]Table 1 The main composition of EVs^[33]

存在部位	组分	组分举例	
		外泌体	微囊泡体
囊泡表面	黏附分子	整合素, 乳凝集素	整合素
	抗原呈递分子	MHC I / II	MHC I / II
	膜运输/融合蛋白	膜联蛋白, 筏蛋白, RAB27	RAB11
	跨膜蛋白	Lamp2b, CD63, CD81	CD63, CD81
	标志蛋白	CD63, CD9	ARF6
囊泡内腔	信号转导蛋白	同线蛋白, G蛋白	ARF6
	骨架蛋白	肌动蛋白, 微管蛋白	肌动蛋白, 微管蛋白
	ESCRT蛋白	ALIX, TSG101, VPS4	ALIX, TSG101
	胞质蛋白	HSP70/90	组蛋白
	酶	氧化物酶, GAPDH	GAPDH
	脂质	鞘磷脂, 胆固醇, 磷脂酰丝氨酸, 神经酰胺	
	核酸	DNA, mRNA, miRNA, ncRNA	

mRNA, miRNA, lncRNA和circRNA), 以及数千种脂质。表1是对其主要成分的总结^[33]。

EVs的脂质成分主要包括鞘磷脂、胆固醇、磷脂酰丝氨酸和神经酰胺, 由于EVs的特殊形成机理, 其主要成分与供体细胞存在很大差别。而EVs中的蛋白质主要包括质膜蛋白和与其形成相关的蛋白, 如膜转运相关蛋白和ESCRT蛋白等。其他还有溶酶体相关膜糖蛋白2b (lysosomal associated membrane glycoprotein 2b, Lamp2b)、四跨膜蛋白、主要组织相容性复合体 I / II (major histocompatibility complex I / II, MHC I / II) 以及甘油醛-3-磷酸脱氢酶(glyceraldehyde-3-phosphate dehydrogenase, GAPDH)等多种蛋白。EVs中也包含了大量的核酸, 其中一些以与蛋白质结合成复合物的形式存在^[35,36]。通过不同的分选机理, 一些特异性的DNA, mRNA, miRNA以及非编码RNA (non-coding RNA, ncRNA)会分选到EVs中, 而且表现出不同的生物学功能。Dvorak等人^[37]首次证明了EVs可以传输完整的mRNA到受体细胞, 并在受体细胞中表达出相应蛋白。EVs中miRNA和长链非编码RNA (long non-coding RNA, lncRNA)也具有重要功能, 如促进肿瘤转移、生长及耐药等^[38-41]。此外, EVs中的多种DNA不仅可以在靶细胞中发挥功能, 还能反映肿瘤的突变状况^[42-44], 如线粒体DNA等^[45]。

在血小板中发现囊泡后, 越来越多的研究表明EVs不仅不是细胞释放的垃圾, 还具有重要的生物学功能,

主要体现在维持细胞生理稳态和参与细胞间信息交流等^[46]。在正常细胞中, EVs参与许多生理过程, 如凝血^[47]、免疫调节^[48]、干细胞分化^[49]、组织再生^[50]和自噬^[51]等。研究表明, 在网织红细胞的成熟过程中, 其还可以通过EVs消除细胞质中的有害DNA和转移受体来避免正常细胞的衰老或凋亡^[52,53]。在肿瘤细胞中, EVs可以作为一种重要的信息传递工具, 与附近甚至远处的细胞传递重要信息, 以建立有利的肿瘤生长和转移条件。肿瘤细胞衍生的EVs对细胞的影响主要包括引发炎症反应、血管生成、细胞迁移、细胞增殖、免疫抑制、肿瘤侵袭和药物抵抗等^[12]。例如, 肿瘤细胞来源的外泌体可以携带特定的RNA, 以激活肺泡上皮细胞的Toll样受体3 (Toll-like receptor 3, TLR3)来招募中性粒细胞, 从而促进肺转移灶的形成^[54]。肿瘤细胞来源的外泌体也可以通过携带不同的整合素来指导癌细胞的转移位置^[55]。而肝癌细胞来源的外泌体还具有活化癌症相关成纤维细胞(cancer-associated fibroblasts, CAFs)的作用, 进而可以促进肝癌细胞向肺的转移^[40]。肿瘤来源的外泌体还可以向树突状细胞(dendritic cells, DCs)传递肿瘤抗原, 以诱导CD8⁺ T细胞依赖性的抗肿瘤作用^[56]。在慢性淋巴白血病中, 外泌体中的RNA可以激活Toll样受体信号通路来活化单核细胞, 从而使单核细胞分泌大量细胞因子, 以重建有利的肿瘤微环境^[57]。因此, EVs无论在正常细胞还是肿瘤细胞中都具有重要生物学功能。

2 EVs的分离和表征

2.1 EVs的分离方法

如上所述, EVs是一类在生物起源、大小及成分上都具有异质性的脂质囊泡, 不同形成机理和成分都可能使EVs具有不同的生物学功能^[58]. 然而, EVs存在于大多数生物体液中, 而这些生物体液中的一些复杂成分, 如核酸、蛋白质及脂质等, 都会影响其功能分析. 因此, 研究EVs生物学功能时, 需要将其从生物体液中分离出来. 目前, 已发展多种分离纯化手段, 包括差速离心、密度梯度离心、免疫捕获、超滤、尺寸排除色谱以及微流控等. 这些方法主要是依靠EVs的大小、密度、形状、电荷和表面成分来实现分离^[59,60].

差速离心最为常用, 是通过不同的离心速度来分离具有不同沉降系数的EVs^[61]. 此方法的优点是可以分离出比较纯的EVs. 但是, EVs的沉降系数会受其形状和密度的影响, 以及EVs会和一些蛋白质、脂质、病毒等产生共沉降, 因此该方法也难以获得高纯度的EVs. 而且高的离心速度还会使EVs产生聚集, 破坏EVs完整性.

密度梯度离心是一种基于大小和密度来分离EVs的方法^[62], 蔗糖和碘克沙醇是常用密度介质. 该方法的优点是可以适当避免蛋白质污染, 分离出高纯度的EVs, 但产量低, 且比较耗时.

免疫捕获是基于微珠或平板表面修饰EVs标志物抗体来捕获EVs的方法, 常用的标志物有CD63和CD81^[63]. 目前, EVs的分离主要是依靠物理学性质(大小和密度). 本方法通过抗体-抗原识别, 可针对含有相关蛋白(抗原)的EVs亚群进行特异性分离. 因此, 更适合用于研究EVs的异质性及功能多样性. 但其分离效率低、成本高, 微珠的结合也会影响EVs的下游分析.

尺寸排阻色谱法是一种基于尺寸大小来分离EVs的方法^[64]. 当EVs混合物通过载有多孔基质的柱子时, 尺寸大于基质孔径的EVs可绕过孔隙被提前洗脱, 而小于孔径的其他成分则能穿过孔隙被后洗脱. 因此, 洗脱时EVs会与其他成分具有不同的流出时间, 从而可实现对EVs的分离. 该方法可以去除95%以上的杂蛋白, 且耗时短, 不会引起EVs的大量聚集和活性的改变. 但分离出的EVs仍会掺杂一些杂蛋白和细胞碎片.

超滤是用具有不同截留尺寸的半透膜来分离EVs^[65]. 此方法快速有效, 能在短时间内分离大量的样品. 但由于外部压力的使用, 会造成EVs的变形.

沉淀法主要是通过聚乙二醇6000/8000 (PEG-6000/8000)来沉淀EVs^[66]. 此方法虽然简单, 但特异性较差, 分离物常含有较多杂蛋白和脂质.

综上所述, 如何实现EVs的高纯度分离仍然是一个大的挑战. 一些新技术也已被尝试用于EVs的分离, 如微流体芯片技术^[67]. 目前, 研究者可以根据研究目的来选择合适的分离技术.

2.2 EVs的表征技术

EVs的表征指标主要包括EVs形状、大小和内容物^[59,68], 采用的手段包括透射电子显微镜、纳米颗粒跟踪分析和荧光活化细胞分选等^[69].

透射电子显微镜是表征EVs结构和尺寸的标准方法, 其分辨率可达0.2 nm左右. 通过电子显微镜成像, 可清楚地看到EVs形态和大小. 纳米颗粒跟踪分析是通过显微镜对粒子布朗运动的跟踪和分析来计算粒子大小和浓度的方法, 检测范围在10~2000 nm之间. 虽然其是EVs常用的计数方法, 但它不能区分与EVs尺寸相同的非EVs^[59]. 荧光活化细胞分选能够鉴别EVs表面蛋白, 分辨率一般在300 nm以上^[70]. 因此对于小EVs的检测, 需要将其吸附到大的微球上, 然后通过荧光标记进行检测. 常用的荧光标记方法是通过抗体对EVs表面特异性标志物的识别来实现, 如膜联蛋白(annexin)和CD63等. 免疫印迹和酶联免疫吸附分析也可以定量分析特定的EVs蛋白^[71]. 若要开展EVs蛋白组分析, 需要采用高通量的质谱法. 目前, EVs中已经有几千种蛋白质被识别, 这对EVs功能研究提供了重要参考. 此外, 扫描电子显微镜、原子力显微镜、动态光散射和电阻脉冲传感等也被用于EVs参数表征^[59].

3 EVs在疾病诊疗中的应用

3.1 EVs作为药物运送载体

如前叙, EVs可作为细胞间的通信者, 携带和转移供体细胞中的生物活性分子, 使其跨越细胞内外障碍从一种细胞传递到另一种细胞^[72]. 更重要的是, EVs携带的这些分子可以在靶细胞中发挥相关功能. 因此, EVs可作为一种新型的药物运送载体, 并具有诸多优点^[5]: (1) EVs的脂质密封结构可以维持药物在体内循环时的稳定性; (2) 其特定的脂质成分(例如鞘磷脂和胆固醇)不仅有助于EVs在体内循环时的稳定性, 还可以刺激EVs与细胞质膜之间的融合; (3) 它们的纳米尺

寸和组成特点可以减小单核吞噬系统(mononuclear phagocytic system, MPS)的识别; (4) 它们源于细胞的性质可以减轻对适应性免疫系统激活; (5) EVs拥有固有的细胞和组织靶向特性。

作为药物运送载体, 控制EVs在体内的稳定性、器官靶向性及生理病理学效应(如免疫刺激和炎症反应)是提高药物治疗效果的关键^[73], 而这些都与EVs的细胞来源相关。因此, EVs供体细胞的选择对药物递送效果具有重要影响。目前, 许多细胞衍生的EVs已被用于药物运送, 如肿瘤细胞或正常293T细胞^[74~76]。间充质干细胞(mesenchymal stem cells, MSCs)由于能分泌大量免疫原性低的EVs, 也被用于药物的运送载体^[77,78]。但MSCs来源的EVs可能会产生副作用, 如促进肿瘤生长及血管生成等^[79]。此外, 不成熟DCs衍生的EVs由于容易获取及免疫原性低等特点, 也是一种潜在的药物运送载体^[80]。随着EVs作为药物运送载体在临床应用中的发展, 其高产率和低成本也越来越重要。现今, 研究者已经开始关注其他来源的EVs或EVs类似物, 如通过微孔和纳米孔连续挤压细胞产生的模拟EVs^[81]。与天然EVs相比, 模拟EVs对细胞中货物的选择是随机的, 缺

乏特异性^[82]。但是, 模拟EVs也已被成功用于药物的递送。通过连续挤压巨噬细胞, 一种包裹阿霉素的模拟EVs在体外和体内都表现出了对肿瘤的抑制作用。同时, 与天然的外泌体载体相比, 模拟EVs具有相似疗效, 且无显著毒性^[83]。

EVs对药物的装载量也是影响治疗效果的关键。目前, 药物装载入EVs主要有两种方法, 即间接载药和直接载药^[84]。

间接载药主要是把细胞内的药物装载到EVs中, 而细胞内的药物可以是自身的, 也可以是额外加入的。为此, 可通过胞内表达核酸(siRNAs, miRNAs, mRNA)或蛋白质的表达来实现^[85~87], 小分子药物等也可以通过电转、转染试剂以及简单的孵育来实现^[77,88]。例如, 通过孵育的方法, DCs来源的EVs不仅实现了对紫杉醇的包装, 还通过携带紫杉醇抑制了胰腺癌细胞的增殖^[89]。但是, 这种间接载药方式的药物装载量有限, 甚至是细胞中表达量最丰富的miRNA在EVs中的包装效率也仅有1%, 即100个EVs中只含有1个miRNA分子^[90,91]。因此, 药物的高效递送需要提高EVs对药物的装载量。目前, 主要的方法是基于EVs的生物起源来实现其对细胞

图 2 (网络版彩色)光调控细胞内源RNA在外泌体中富集和释放的原理^[94]。该系统由3个主要步骤组成。第一步是RNA的富集。Palm-EGFP-CIBN融合蛋白通过Palm序列固定在质膜上, 然后蓝光介导的可逆CIBN-CRY2相互作用将CRY2-mCherry-MCP招募到质膜上。MCP与MS2的相互作用再使BFP-miR-21sponge-MS2富集于质膜。第二步是工程化外泌体的生物起源和释放。第三步是靶向递送。胆固醇修饰的AS1411适体通过脂质和胆固醇的相互作用固定在外泌体表面, 工程化外泌体再通过AS1411和核仁素的相互作用靶向白血病细胞。

Figure 2 (Color online) The schematic for light regulate the enrichment and release of endogenous RNA in exosomes^[94]. This system consists of three major steps. Step I is the enrichment of RNA. Palm-EGFP-CIBN conjugates are fixed on the plasma membrane by the palm sequence, and then, CRY2-mCherry-MCP is recruited to the plasma membrane by a blue light-mediated reversible CIBN-CRY2 interaction. Next, BFP-miR-21sponge-MS2 also is enriched on the plasma membrane by the interaction of MCP and MS2. Step II is biogenesis and the release of engineered exosomes. Step III is targeted therapy. Cholesterol labeled AS1411 aptamer was attached on the surface of exosomes by the interaction of cholesterol and lipid of exosomes. The engineered exosomes can target leukemia cells through the interaction of AS1411 and nucleolin.

内源蛋白质或RNA的富集^[86,92,93]。为此,我们通过对外泌体供体细胞的设计,发展了一种基于外泌体的新型光控纳米递送系统,实现了内源RNA治疗剂的高效装载及对肿瘤细胞的靶向递送(图2)^[94]。该系统通过RNA适配体(MS2)-蛋白质(MCP)相互作用和光诱导可逆蛋白(CIBN)-蛋白(CRY2)相互作用,实现了活细胞中内源表达miRNA海绵(miRNA21 sponge)在细胞质膜上大量富集。进而,通过外泌体生物起源,miRNA21海绵作为治疗剂被高效装载至工程化外泌体中。最后,通过适配体AS1411对外泌体修饰,装载miR-21海绵的外泌体能够靶向白血病细胞,从而阻断miR-21的功能,诱导白血病细胞凋亡。

直接载药是把药物直接载入分离纯化后的EVs中。其中,电穿孔是最常用的方法。通过该方法,Matthew Wood小组实现了siRNA在EVs中的装载,并使siRNA在阿尔茨海默病小鼠模型中成功跨越了血脑屏障进行功能传递^[95]。而其他研究小组还成功地将1000 bp的DNA^[96]、5 nm的超顺磁性氧化铁纳米粒子和大蛋白质加载到了EVs中^[97]。然而,电穿孔法可能诱导siRNA和EVs的共聚集,从而产生过高估计的实际装载效率^[98]。转染也是一种常用的EVs载药法。例如,siRNA通过转染被成功加载到EVs后,也实现了对乳腺癌的抑制^[99]。反复冻融循环、超声、挤压或皂素等处理也可用于EVs的药物装载^[100],但这些方法都难以保证EVs的完整性。对于疏水膜渗透性分子,简单的孵育可实现EVs对药物的装载并保证EVs的完整性。该方法使EVs携带了姜黄素到GL26小鼠肿瘤模型,抑制了脑瘤的生长^[101]。

目前,EVs虽然已被成功用于多种治疗性药物的递送载体。然而,EVs作为药物递送载体应用于临床前,还需要对EVs的分离方法和药物装载技术进一步改进,以及更深层次研究EVs组成、异质性、递送机制和生物学效应等。

3.2 EVs作为疾病诊断标志物

基于EVs在体液中稳定存在及与母细胞具有相似成分的特点,EVs有望成为肿瘤早期诊断和液体活检的生物标志物^[102]。虽然,对EVs成分的调控机理还未完全了解,但大量研究已经证明EVs中存在与肿瘤相关的蛋白质和核酸^[103~106]。相对于组织活检,基于EVs的液体活检不仅提供了一种更方便和非侵入性的诊断手段,还能更好体现肿瘤异质性及肿瘤的发生发展^[107]。

EVs中肿瘤标志物的发现主要基于质谱分析、

RNA测序、蛋白质芯片和核酸芯片等技术。例如,质谱分析发现,磷脂酰肌醇蛋白聚糖-1 (glypican-1, GPC1)阳性的EVs水平可作为一种潜在的非侵入性诊断和筛查工具,用于早期胰腺癌的检测^[69]。纳米等离子体共振技术对卵巢癌病人EVs的分析发现,CD24和上皮细胞黏附分子(epithelial cell adhesion molecules, EpCAM)在其上存在差异性表达,表明其可能具有作为诊断标志物的潜力^[108,109]。在乳腺癌患者中,免疫印迹分析证明,EVs上的生存素(survivin)水平明显高于对照组,表明生存素阳性的EVs可能来源于肿瘤细胞^[110]。通过质谱分析,我们发现核仁素可能是一种白血病相关外泌体上的标志物之一。为此,我们发展了一种针对核仁素和CD63的外泌体双信号放大检测方法(图3),检测灵敏度达 $1 \times 10^2/\mu\text{L}$,为白血病早期诊断、液体活检提供了新的选择^[111]。除了蛋白质,EVs中的miRNA也具有作为肿瘤诊断标志物的潜力^[112]。

越来越多的研究证明,EVs不仅可以用于肿瘤诊断,还可以用于预测肿瘤的发生发展及临床用药结果^[113]。但在广泛应用于临床之前,还有许多问题需要解决。例如,尚未在EVs中发现高特异性的肿瘤标志物,EVs标准纯化方案的缺乏制约了对生物标志物识别的再现性和重复性^[5]、低丰度肿瘤EVs检出率低等^[114]。

3.3 EVs作为治疗剂

基于EVs可携带供体细胞生物活性成分的特殊性质,DCs衍生的EVs被首次作为一种无细胞抗癌疫苗,为EVs提供了第一个治疗应用^[115]。随后,DCs衍生的EVs还被用于非小细胞肺癌^[116,117]和黑色素瘤的疫苗^[118],并分别进入了临床一期和二期阶段。此外,抗原呈递细胞(antigen-presenting cells, APCs)衍生的EVs可以通过呈递MHC到T细胞,进而刺激免疫系统^[119]。除了免疫细胞,MSCs衍生的EVs也可作为自身细胞的代替物用于肿瘤治疗。例如,在自体免疫性疾病1型糖尿病中,基于MSCs衍生EVs的治疗方案已经进入临床二期^[120]。而且,研究表明MSCs衍生的EVs还具有促进组织修复和再生的能力^[121]。虽然MSCs衍生EVs的治疗机理尚不明确,但EVs中包含的很多细胞因子,如趋化因子,白细胞介素和生长因子等信号蛋白,可能发挥了重要作用。除上述几种,其他细胞衍生的EVs也被报道可以用于肿瘤的治疗,如自然杀伤细胞(natural killer cells, NK)衍生的EVs可以通过携带穿孔蛋白(perforin)引起癌细胞死亡^[122],对黑色素瘤细胞产生毒性等^[123]。

图 3 (网络版彩色)外泌体双靶标分子(核仁素和CD63)的双信号放大检测原理^[111]。反应步骤为: (1) 用抗CD63抗体修饰磁珠偶联物(MB-CD63)捕获白血病细胞来源外泌体; (2) 使用包含核仁素蛋白适体(AS1411)的DNA引物与外泌体结合; (3) 启动滚环扩增反应(Rolling circle amplification, RCA); (4) 与金纳米颗粒(GNP)-DNA-荧光染料(FAM)偶联物(GNP-DNA-FAM)杂交; (5) 通过核酸内切酶(Nb.BbvCI)辅助的靶标循环, 使FAM从GNP-DNA-FAM偶联物中释放出来, 由猝灭状态转化为发射状态, 荧光信号不断积累

Figure 3 (Color online) Principle of double signal amplification detection of exosome by two target molecules (nucleolin and CD63)^[111]. First, leukemia-derived exosomes were captured by anti-CD63 antibody modified magnetic beads conjugates (MB-CD63). Second, primer containing aptamer of nucleolin (AS1411) was applied to bind exosomes. Third, initiating the rolling circle amplification (RCA). Fourth, the hybridization of RCA product with GNP-DNA-FAM conjugates. Finally, nicking endonuclease (Nb.BbvCI) assisted target recycling was introduced. As a result, FAM was released from GNP-DNA-FAM conjugates, transforming FAM from quenching state to emission state and thus fluorescence signal continuously accumulated

如上所述, EVs作为治疗剂具有很大的潜力, 已被广泛研究. 与其母细胞相比, EVs易储存, 生物安全性也更好. 但是, EVs作为治疗剂, 机理尚不清楚, 而且其定义、标准化, 以及安全性评估均需研究.

4 结语

EVs在生物体系内如此广泛存在, 其生物学功能也越来越多地被认识, 或许正在成为一个“细胞外囊泡生

物学”领域. 对EVs性质的认识也促进了相关应用研究, 涉及疾病诊断、治疗和药物运送等. 然而, 现有EVs的研究方法尚不成熟, 成为阻碍临床转化应用的瓶颈. 对EVs形成和调控机理的深入认识, 一方面有助于我们更好地理解EVs的生物学功能、异质性和功能多样性. 另一方面有可能基于这些知识来发展更有效的EVs改造和制备方法, 并实现工程化和标准化. 随着研究的深入和方法学创新, 这些问题将会逐步得到解决.

参考文献

- 1 Raposo G, Stoorvogel W. Extracellular vesicles: Exosomes, microvesicles, and friends. *J Cell Biol*, 2013, 200: 373–383
- 2 Willms E, Cabañas C, Mäger I, et al. Extracellular vesicle heterogeneity: Subpopulations, isolation techniques, and diverse functions in cancer progression. *Front Immunol*, 2018, 9: 738
- 3 Wolf P. The nature and significance of platelet products in human plasma. *Br J Haematol*, 1967, 13: 269–288
- 4 Pan B T. Electron microscopic evidence for externalization of the transferrin receptor in vesicular form in sheep reticulocytes. *J Cell Biol*, 1985, 101: 942–948
- 5 Stremersch S, De Smedt S C, Raemdonck K. Therapeutic and diagnostic applications of extracellular vesicles. *J Cell Biol*, 2016, 244: 167–183
- 6 Pap E, Pállinger É, Pásztói M, et al. Highlights of a new type of intercellular communication: microvesicle-based information transfer. *Inflamm*

- Res, 2009, 58: 1–8
- 7 Muralidharan-Chari V, Clancy J, Plou C, et al. ARF6-regulated shedding of tumor cell-derived plasma membrane microvesicles. *Curr Biol*, 2009, 19: 1875–1885
- 8 Piccin A, Murphy W G, Smith O P. Circulating microparticles: Pathophysiology and clinical implications. *Blood Rev*, 2007, 21: 157–171
- 9 del Conde I. Tissue-factor-bearing microvesicles arise from lipid rafts and fuse with activated platelets to initiate coagulation. *Blood*, 2005, 106: 1604–1611
- 10 Al-Nedawi K, Meehan B, Micallef J, et al. Erratum. *Nat Cell Biol*, 2008, 10: 752
- 11 Di Vizio D, Kim J, Hager M H, et al. Oncosome formation in prostate cancer: Association with a region of frequent chromosomal deletion in metastatic disease. *Cancer Res*, 2009, 69: 5601–5609
- 12 Minciacchi V R, Freeman M R, Di Vizio D. Extracellular vesicles in cancer: Exosomes, microvesicles and the emerging role of large oncosomes. *Seminars Cell Dev Biol*, 2015, 40: 41–51
- 13 Johnstone R M, Adam M, Hammond J R, et al. Vesicle formation during reticulocyte maturation-association of plasma-membrane activities with released vesicles (exosomes). *J Biol Chem*, 1987, 262: 9412–9420
- 14 Théry C, Zitvogel L, Amigorena S. Exosomes: Composition, biogenesis and function. *Nat Rev Immunol*, 2002, 2: 569–579
- 15 Kowal J, Tkach M, Théry C. Biogenesis and secretion of exosomes. *Curr Opin Cell Biol*, 2014, 29: 116–125
- 16 Gross J C, Chaudhary V, Bartscherer K, et al. Active Wnt proteins are secreted on exosomes. *Nat Cell Biol*, 2012, 14: 1036–1045
- 17 Tamai K, Tanaka N, Nakano T, et al. Exosome secretion of dendritic cells is regulated by Hrs, an ESCRT-0 protein. *Biochem Biophys Res Commun*, 2010, 399: 384–390
- 18 Colombo M, Moita C, van Niel G, et al. Analysis of ESCRT functions in exosome biogenesis, composition and secretion highlights the heterogeneity of extracellular vesicles. *J Cell Sci*, 2013, 126: 5553–5565
- 19 Baietti M F, Zhang Z, Mortier E, et al. Syndecan-syntenin-ALIX regulates the biogenesis of exosomes. *Nat Cell Biol*, 2012, 14: 677–685
- 20 Perez-Hernandez D, Gutiérrez-Vázquez C, Jorge I, et al. The intracellular interactome of tetraspanin-enriched microdomains reveals their function as sorting machineries toward exosomes. *J Biol Chem*, 2013, 288: 11649–11661
- 21 Verweij F J, van Eijndhoven M A J, Hopmans E S, et al. LMP1 association with CD63 in endosomes and secretion via exosomes limits constitutive NF- κ B activation. *EMBO J*, 2011, 30: 2115–2129
- 22 Trajkovic K. Ceramide triggers budding of exosome vesicles into multivesicular endosomes. *Science*, 2008, 320: 179
- 23 Laulagnier K, Grand D, Dujardin A, et al. PLD2 is enriched on exosomes and its activity is correlated to the release of exosomes. *FEBS Lett*, 2004, 572: 11–14
- 24 Sahu R, Kaushik S, Clement C C, et al. Microautophagy of cytosolic proteins by late endosomes. *Dev Cell*, 2011, 20: 405–406
- 25 Mittelbrunn M, Gutiérrez-Vázquez C, Villarroya-Beltri C, et al. Unidirectional transfer of microRNA-loaded exosomes from T cells to antigen-presenting cells. *Nat Commun*, 2011, 2: 282
- 26 Villarroya-Beltri C, Baixauli F, Gutiérrez-Vázquez C, et al. Sorting it out: Regulation of exosome loading. *Seminars Cancer Biol*, 2014, 28: 3–13
- 27 Sinha S, Hoshino D, Hong N H, et al. Cortactin promotes exosome secretion by controlling branched actin dynamics. *J Cell Biol*, 2016, 214: 197–213
- 28 Stenmark H. Rab GTPases as coordinators of vesicle traffic. *Nat Rev Mol Cell Biol*, 2009, 10: 513–525
- 29 Bobrie A, Krumeich S, Reyat F, et al. Rab27a supports exosome-dependent and -independent mechanisms that modify the tumor microenvironment and can promote tumor progression. *Cancer Res*, 2012, 72: 4920–4930
- 30 Peinado H, Alečković M, Lavotshkin S, et al. Melanoma exosomes educate bone marrow progenitor cells toward a pro-metastatic phenotype through MET. *Nat Med*, 2012, 18: 883–891
- 31 Ostrowski M, Carmo N B, Krumeich S, et al. Rab27a and Rab27b control different steps of the exosome secretion pathway. *Nat Cell Biol*, 2010, 12: 19–30
- 32 Hsu C, Morohashi Y, Yoshimura S, et al. Regulation of exosome secretion by Rab35 and its GTPase-activating proteins TBC1D10A–C. *J Cell Biol*, 2010, 189: 223–232
- 33 Colombo M, Raposo G, Théry C. Biogenesis, secretion, and intercellular interactions of exosomes and other extracellular vesicles. *Annu Rev Cell Dev Biol*, 2014, 30: 255–289
- 34 Abels E R, Breakefield X O. Introduction to extracellular vesicles: Biogenesis, RNA cargo selection, content, release, and uptake. *Cell Mol Neurobiol*, 2016, 36: 301–312
- 35 Villarroya-Beltri C, Gutiérrez-Vázquez C, Sánchez-Cabo F, et al. Sumoylated hnRNPA2B1 controls the sorting of miRNAs into exosomes through binding to specific motifs. *Nat Commun*, 2013, 4: 2980
- 36 Kossinova O A, Gopanenko A V, Tamkovich S N, et al. Cytosolic YB-1 and NSUN2 are the only proteins recognizing specific motifs present in mRNAs enriched in exosomes. *Biochim Biophys Acta-Proteins Proteom*, 2017, 1865: 664–673

- 37 Ratajczak J, Miekus K, Kucia M, et al. Embryonic stem cell-derived microvesicles reprogram hematopoietic progenitors: Evidence for horizontal transfer of mRNA and protein delivery. *Leukemia*, 2006, 20: 847–856
- 38 Ying W, Riopel M, Bandyopadhyay G, et al. Adipose tissue macrophage-derived exosomal mirnas can modulate *in vivo* and *in vitro* insulin sensitivity. *Cell*, 2017, 171: 372–384.e12
- 39 Yan W, Wu X, Zhou W, et al. Cancer-cell-secreted exosomal miR-105 promotes tumour growth through the MYC-dependent metabolic reprogramming of stromal cells. *Nat Cell Biol*, 2018, 20: 597–609
- 40 Fang T, Lv H, Lv G, et al. Tumor-derived exosomal miR-1247-3p induces cancer-associated fibroblast activation to foster lung metastasis of liver cancer. *Nat Commun*, 2018, 9: 191
- 41 Qu L, Ding J, Chen C, et al. Exosome-transmitted lncARSR promotes sunitinib resistance in renal cancer by acting as a competing endogenous RNA. *Cancer Cell*, 2016, 29: 653–668
- 42 Balaj L, Lessard R, Dai L, et al. Tumour microvesicles contain retrotransposon elements and amplified oncogene sequences. *Nat Commun*, 2011, 2: 180
- 43 Thakur B K, Zhang H, Becker A, et al. Double-stranded DNA in exosomes: A novel biomarker in cancer detection. *Cell Res*, 2014, 24: 766–769
- 44 Lázaro-Ibáñez E, Sanz-García A, Visakorpi T, et al. Different gDNA content in the subpopulations of prostate cancer extracellular vesicles: Apoptotic bodies, microvesicles, and exosomes. *Prostate*, 2014, 74: 1379–1390
- 45 Guescini M, Genedani S, Stocchi V, et al. Astrocytes and glioblastoma cells release exosomes carrying mtDNA. *J Neural Transm*, 2010, 117: 1–4
- 46 Budnik V, Ruiz-Cañada C, Wendler F. Extracellular vesicles round off communication in the nervous system. *Nat Rev Neurosci*, 2016, 17: 160–172
- 47 Heijnen H F G, Schiel A E, Fijnheer R, et al. Activated platelets release two types of membrane vesicles: Microvesicles by surface shedding and exosomes derived from exocytosis of multivesicular bodies and alpha-granules. *Blood*, 1999, 94: 3791–3799
- 48 Greening D W, Gopal S K, Xu R, et al. Exosomes and their roles in immune regulation and cancer. *Seminars Cell Dev Biol*, 2015, 40: 72–81
- 49 Nair R, Santos L, Awasthi S, et al. Extracellular vesicles derived from preosteoblasts influence embryonic stem cell differentiation. *Stem Cells Dev*, 2014, 23: 1625–1635
- 50 Teng X, Chen L, Chen W, et al. Mesenchymal stem cell-derived exosomes improve the microenvironment of infarcted myocardium contributing to angiogenesis and anti-inflammation. *Cell Physiol Biochem*, 2015, 37: 2415–2424
- 51 Xu J, Camfield R, Gorski S M. The interplay between exosomes and autophagy-partners in crime. *J Cell Sci*, 2018, 131: jcs215210
- 52 Takahashi A, Okada R, Nagao K, et al. Exosomes maintain cellular homeostasis by excreting harmful DNA from cells. *Nat Commun*, 2017, 8: 15287
- 53 Pan B T, Johnstone R M. Fate of the transferrin receptor during maturation of sheep reticulocytes *in vitro*: Selective externalization of the receptor. *Cell*, 1983, 33: 967–978
- 54 Liu Y, Gu Y, Han Y, et al. Tumor exosomal rnas promote lung pre-metastatic niche formation by activating alveolar epithelial TLR3 to recruit neutrophils. *Cancer Cell*, 2016, 30: 243–256
- 55 Hoshino A, Costa-Silva B, Shen T L, et al. Tumour exosome integrins determine organotropic metastasis. *Nature*, 2015, 527: 329–335
- 56 Wolfers J, Lozier A, Raposo G, et al. Tumor-derived exosomes are a source of shared tumor rejection antigens for CTL cross-priming. *Nat Med*, 2001, 7: 297–303
- 57 Haderk F, Schulz R, Iskar M, et al. Tumor-derived exosomes modulate PD-L1 expression in monocytes. *Sci Immunol*, 2017, 2: eaah5509
- 58 Théry C, Ostrowski M, Segura E. Membrane vesicles as conveyors of immune responses. *Nat Rev Immunol*, 2009, 9: 581
- 59 Coumans F A W, Brisson A R, Buzas E I, et al. Methodological guidelines to study extracellular vesicles. *Circ Res*, 2017, 120: 1632–1648
- 60 Marques-García F, Isidoro-García M. Protocols for exosome isolation and RNA profiling. *Methods Mol Biol*, 2016, 1434: 153–167
- 61 Valadi H, Ekström K, Bossios A, et al. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol*, 2007, 9: 654–659
- 62 Thery C, Amigorena S, Raposo G, et al. Isolation and characterization of exosomes from cell culture supernatants and biological fluids. *Curr Protoc Cell Biol*, 2006, Chapter 3: Unit 3 22, doi: 10.1002/0471143030.cb0322s30
- 63 Oksvold M P, Neurauter A, Pedersen K W. Magnetic bead-based isolation of exosomes. *Methods Mol Biol*, 2015, 1218: 465–481
- 64 Merchant M L, Powell D W, Wilkey D W, et al. Microfiltration isolation of human urinary exosomes for characterization by MS. *Prot Clin Appl*, 2010, 4: 84–96
- 65 Alvarez M L, Khosroheidari M, Kanchi R R, et al. Comparison of protein, microRNA, and mRNA yields using different methods of urinary exosome isolation for the discovery of kidney disease biomarkers. *Kidney Int*, 2012, 82: 1024–1032
- 66 Lane R E, Korbie D, Anderson W, et al. Analysis of exosome purification methods using a model liposome system and tunable-resistive pulse sensing. *Sci Rep*, 2015, 5: 7639
- 67 Liu C, Guo J, Tian F, et al. Field-free isolation of exosomes from extracellular vesicles by microfluidic viscoelastic flows. *ACS Nano*, 2017, 11:

6968–6976

- 68 Shao H, Im H, Castro C M, et al. New technologies for analysis of extracellular vesicles. *Chem Rev*, 2018, 118: 1917–1950
- 69 Melo S A, Luecke L B, Kahlert C, et al. Glypican-1 identifies cancer exosomes and detects early pancreatic cancer. *Nature*, 2015, 523: 177–182
- 70 Shen W, Guo K, Adkins G B, et al. A single extracellular vesicle (EV) flow cytometry approach to reveal EV heterogeneity. *Angew Chem Int Ed*, 2018, 57: 15675–15680
- 71 Kim Y, Jeon J, Mejia S, et al. Targeted proteomics identifies liquid-biopsy signatures for extracapsular prostate cancer. *Nat Commun*, 2016, 7: 11906
- 72 Tkach M, Théry C. Communication by extracellular vesicles: Where we are and where we need to go. *Cell*, 2016, 164: 1226–1232
- 73 Johnsen K B, Gudbergsson J M, Skov M N, et al. A comprehensive overview of exosomes as drug delivery vehicles-endogenous nanocarriers for targeted cancer therapy. *Biochim Biophys Acta-Rev Cancer*, 2014, 1846: 75–87
- 74 Maguire C A, Balaj L, Sivaraman S, et al. Microvesicle-associated AAV vector as a novel gene delivery system. *Mol Ther*, 2012, 20: 960–971
- 75 Shtam T A, Kovalev R A, Varfolomeeva E, et al. Exosomes are natural carriers of exogenous siRNA to human cells *in vitro*. *Cell Commun Signal*, 2013, 11: 88
- 76 Aspe J R, Diaz Osterman C J, Jutzy J M S, et al. Enhancement of Gemcitabine sensitivity in pancreatic adenocarcinoma by novel exosome-mediated delivery of the Survivin-T34A mutant. *J ExtraCellular Vesicles*, 2014, 3: 23244
- 77 Munoz J L, Bliss S A, Greco S J, et al. Delivery of functional anti-miR-9 by mesenchymal stem cell-derived exosomes to glioblastoma multiforme cells conferred chemosensitivity. *Mol Ther-Nucleic Acids*, 2013, 2: e126
- 78 Xin H, Li Y, Buller B, et al. Exosome-mediated transfer of miR-133b from multipotent mesenchymal stromal cells to neural cells contributes to neurite outgrowth. *Stem Cells*, 2012, 30: 1556–1564
- 79 Zhu W, Huang L, Li Y, et al. Exosomes derived from human bone marrow mesenchymal stem cells promote tumor growth *in vivo*. *Cancer Lett*, 2012, 315: 28–37
- 80 Yin W, Ouyang S, Li Y, et al. Immature dendritic cell-derived exosomes: A promise subcellular vaccine for autoimmunity. *Inflammation*, 2013, 36: 232–240
- 81 Yang Z, Xie J, Zhu J, et al. Functional exosome-mimic for delivery of siRNA to cancer: *In vitro* and *in vivo* evaluation. *J Control Release*, 2016, 243: 160–171
- 82 García-Manrique P, Gutiérrez G, Blanco-López M C. Fully artificial exosomes: Towards new theranostic biomaterials. *Trends Biotech*, 2018, 36: 10–14
- 83 Jang S C, Kim O Y, Yoon C M, et al. Bioinspired exosome-mimetic nanovesicles for targeted delivery of chemotherapeutics to malignant tumors. *ACS Nano*, 2013, 7: 7698–7710
- 84 Piffoux M, Nicolas-Boluda A, Mulens-Arias V, et al. Extracellular vesicles for personalized medicine: The input of physically triggered production, loading and theranostic properties. *Adv Drug Deliv Rev*, 2019, 138: 247–258
- 85 Kosaka N, Iguchi H, Yoshioka Y, et al. Secretory mechanisms and intercellular transfer of microRNAs in living cells. *J Biol Chem*, 2010, 285: 17442–17452
- 86 Yim N, Ryu S W, Choi K, et al. Exosome engineering for efficient intracellular delivery of soluble proteins using optically reversible protein-protein interaction module. *Nat Commun*, 2016, 7: 12277
- 87 Bolukbasi M F, Mizrak A, Ozdener G B, et al. miR-1289 and “zipcode”-like sequence enrich mRNAs in microvesicles. *Mol Ther-Nucleic Acids*, 2012, 1: e10
- 88 Ohno S, Takanashi M, Sudo K, et al. Systemically injected exosomes targeted to EGFR deliver antitumor microrna to breast cancer cells. *Mol Ther*, 2013, 21: 185–191
- 89 Pascucci L, Coccè V, Bonomi A, et al. Paclitaxel is incorporated by mesenchymal stromal cells and released in exosomes that inhibit *in vitro* tumor growth: A new approach for drug delivery. *J Control Release*, 2014, 192: 262–270
- 90 Tosar J P, Gambaro F, Sanguinetti J, et al. Assessment of small RNA sorting into different extracellular fractions revealed by high-throughput sequencing of breast cell lines. *Nucleic Acids Res*, 2015, 43: 5601–5616
- 91 Chevillet J R, Kang Q, Ruf I K, et al. Quantitative and stoichiometric analysis of the microRNA content of exosomes. *Proc Natl Acad Sci USA*, 2014, 111: 14888–14893
- 92 Wang Q, Yu J, Kadungure T, et al. ARMMs as a versatile platform for intracellular delivery of macromolecules. *Nat Commun*, 2018, 9: 960
- 93 Kojima R, Bojar D, Rizzi G, et al. Designer exosomes produced by implanted cells intracerebrally deliver therapeutic cargo for Parkinson’s disease treatment. *Nat Commun*, 2018, 9: 1305
- 94 Huang L, Gu N, Zhang X E, et al. Light-inducible exosome-based vehicle for endogenous RNA loading and delivery to leukemia cells. *Adv Funct Mater*, 2019: 1807189
- 95 Alvarez-Erviti L, Seow Y, Yin H F, et al. Delivery of siRNA to the mouse brain by systemic injection of targeted exosomes. *Nat Biotechnol*, 2011,

29: 341–345

- 96 Lamichhane T N, Raiker R S, Jay S M. Exogenous DNA loading into extracellular vesicles via electroporation is size-dependent and enables limited gene delivery. *Mol Pharm*, 2015, 12: 3650–3657
- 97 Hood J L, Scott M J, Wickline S A. Maximizing exosome colloidal stability following electroporation. *Anal Biochem*, 2014, 448: 41–49
- 98 Kooijmans S A A, Stremersch S, Braeckmans K, et al. Electroporation-induced siRNA precipitation obscures the efficiency of siRNA loading into extracellular vesicles. *J Control Release*, 2013, 172: 229–238
- 99 Pi F, Binzel D W, Lee T J, et al. Nanoparticle orientation to control RNA loading and ligand display on extracellular vesicles for cancer regression. *Nat Nanotech*, 2018, 13: 82–89
- 100 Haney M J, Klyachko N L, Zhao Y, et al. Exosomes as drug delivery vehicles for Parkinson's disease therapy. *J Control Release*, 2015, 207: 18–30
- 101 Zhuang X, Zhang H G. Abstract 4831: Treatment of brain inflammatory diseases by delivering exosome encapsulated anti-inflammatory drugs from the nasal region to the brain. *Cancer Res*, 2012, 72: 4831
- 102 Soung Y, Ford S, Zhang V, et al. Exosomes in cancer diagnostics. *Cancers*, 2017, 9: 8
- 103 Caivano A, Laurenzana I, De Luca L, et al. High serum levels of extracellular vesicles expressing malignancy-related markers are released in patients with various types of hematological neoplastic disorders. *Tumor Biol*, 2015, 36: 9739–9752
- 104 De Luca L, D'Arena G, Simeon V, et al. Characterization and prognostic relevance of circulating microvesicles in chronic lymphocytic leukemia. *Leukemia Lymphoma*, 2017, 58: 1424–1432
- 105 Shin K J, Seo Y K, Yoo Y. Proteomic profiling of exosomes leads to the identification of novel biomarkers for prostate cancer progression. *Faseb J*, 2018, 32: e82589
- 106 Momen-Heravi F, Getting S J, Moschos S A. Extracellular vesicles and their nucleic acids for biomarker discovery. *Pharmacol Therapeut*, 2018, 192: 170–187
- 107 Hofman P. Liquid biopsy for early detection of lung cancer. *Curr Opin Oncol*, 2017, 29: 73–78
- 108 Im H, Shao H, Park Y I, et al. Label-free detection and molecular profiling of exosomes with a nano-plasmonic sensor. *Nat Biotechnol*, 2014, 32: 490–495
- 109 Zhao Z, Yang Y, Zeng Y, et al. A microfluidic ExoSearch chip for multiplexed exosome detection towards blood-based ovarian cancer diagnosis. *Lab Chip*, 2016, 16: 489–496
- 110 Khan S, Bennit H F, Turay D, et al. Early diagnostic value of survivin and its alternative splice variants in breast cancer. *BMC Cancer*, 2014, 14: 176
- 111 Huang L, Wang D B, Singh N, et al. A dual-signal amplification platform for sensitive fluorescence biosensing of leukemia-derived exosomes. *Nanoscale*, 2018, 10: 20289–20295
- 112 Rabinowits G, Gerçel-Taylor C, Day J M, et al. Exosomal microRNA: A diagnostic marker for lung cancer. *Clin Lung Cancer*, 2009, 10: 42–46
- 113 Chen G, Huang A C, Zhang W, et al. Exosomal PD-L1 contributes to immunosuppression and is associated with anti-PD-1 response. *Nature*, 2018, 560: 382–386
- 114 Xu R, Rai A, Chen M, et al. Extracellular vesicles in cancer—Implications for future improvements in cancer care. *Nat Rev Clin Oncol*, 2018, 15: 617–638
- 115 Zitvogel L, Regnault A, Lozier A, et al. Eradication of established murine tumors using a novel cell-free vaccine: Dendritic cell derived exosomes. *Nat Med*, 1998, 4: 594–600
- 116 Morse M A, Garst J, Osada T, et al. A phase I study of dexosome immunotherapy in patients with advanced non-small cell lung cancer. *J Transl Med*, 2005, 3: 9
- 117 Viaud S, Ploix S, Lapierre V, et al. Updated technology to produce highly immunogenic dendritic cell-derived exosomes of clinical grade. *J Immunother*, 2011, 34: 65–75
- 118 Escudier B, Dorval T, Chaput N, et al. Vaccination of metastatic melanoma patients with autologous dendritic cell (DC) derived-exosomes: Results of the first phase I clinical trial. *J Transl Med*, 2005, 3: 10
- 119 Raposo G, Nijman H W, Stoorvogel W, et al. B-lymphocytes secrete antigen-presenting vesicles. *Mol Biol Cell*, 1995, 6: 1248
- 120 Favaro E, Carpanetto A, Lamorte S, et al. Human mesenchymal stem cell-derived microvesicles modulate T cell response to islet antigen glutamic acid decarboxylase in patients with type 1 diabetes. *Diabetologia*, 2014, 57: 1664–1673
- 121 Garikipati V N S, Shoja-Taheri F, Davis M E, et al. Extracellular vesicles and the application of system biology and computational modeling in cardiac repair. *Circ Res*, 2018, 123: 188–204
- 122 Lugini L, Cecchetti S, Huber V, et al. Immune surveillance properties of human nk cell-derived exosomes. *J Immunol*, 2012, 189: 2833–2842
- 123 Zhu L, Kalimuthu S, Gangadaran P, et al. Exosomes derived from natural killer cells exert therapeutic effect in melanoma. *Theranostics*, 2017, 7: 2732–2745

Summary for “细胞外囊泡及其在疾病诊疗中的应用”

Extracellular vesicles and their application in the diagnosis and treatment of diseases

Lin Huang^{1,2†}, Dianbing Wang^{2†}, Ning Gu¹ & Xian-en Zhang^{1,2*}¹ School of Biological Science and Medical Engineering, Southeast University, Nanjing 210096, China;² National Laboratory of Biomacromolecules, CAS Center for Excellence in Biomacromolecules, Institute of Biophysics, Chinese Academy of Sciences, Beijing 100101, China

† Equally contributed to this work.

* Corresponding author, E-mail: zhangxe@ibp.ac.cn

Extracellular vesicles (EVs) are a kind of heterogeneous vesicles released by essentially all cell types, which plays an important role in intercellular communication and in the regulation of physiological process (or pathology). EVs mainly includes microvesicles (50–1000 nm), apoptotic bodies (50–5000 nm) and exosomes (30–150 nm), based on distinct biogenesis pathways. Microvesicles are formed by direct budding of the plasma membrane, in which ARF6 and RhoA are important regulators. In addition, oncosomes are also a kind of microvesicles, which are specifically released by tumor cells and contain many specific molecules of tumor cells. When cells undergo apoptosis, they could also release a population of EVs called apoptotic bodies. This kind of EVs is formed through outward blebbing and fragmentation of the cell membrane. Exosomes are derived from multivesicular bodies (MVBs), and released by cells upon fusion of MVBs with the plasma membrane. Exosomes have complex formation and regulatory mechanisms, including endosomal sorting complex dependent and non-dependent mechanisms.

Many researchers have shown that EVs with different types and different cell sources may have different components and functions. EVs mainly carry the bioactive molecules of donor cells, such as proteins, DNA or RNA, and target nearby or remote cells to perform different biological functions, thus affecting the physiological and pathological processes of recipient cells. Cells can also discharge their waste by EVs. By transporting cargoes to extracellular, EVs can regulate the amount of intracellular cargoes, thus affecting the survival status of cells. This approach can be beneficial to cells (removing harmful substances) or harmful (transporting drugs outside the cell). Secondly, some key molecules in EVs can be regulated by donor cells according to their state. So, EVs can also reflect the state changes of donor cells, and have the potential as a biomarker for the diagnosis and treatment of diseases. Due to their stable existence in body fluids and derived from various tissues of the body, EVs can not only be a new biomarker of liquid biopsy, but also reflect the heterogeneity and the genesis or development of diseases. As a natural endogenous material, EVs can also serve as a potential drug delivery carrier, and have the advantages of low immunogenicity and crossing the blood brain barrier. In addition, EVs have many other applications. For example, they can be an alternative to cells as a therapeutic agent, as well as targets of disease treatment.

This review summarizes the following parts: Firstly, biological origin, biological composition, biological characteristics, and biological functions of EVs. Secondly, the isolation and characterization of EVs. Thirdly, the application of EVs in the diagnosis and treatment of diseases were introduced. At present, the heterogeneity, biological composition and the formation mechanism of EVs need to be further studied, so that it can be better applied in the diagnosis and treatment of diseases. The effective isolation method is also necessary for its clinical application. It is believed that with the development of various biological technologies, these problems will be gradually solved, the biological function of EVs will be also clearly understood, and EVs based application in diseases will make new breakthroughs.

extracellular vesicles, drug delivery carrier, exosomes, nanomaterials, tumor markerdoi: [10.1360/N972019-00018](https://doi.org/10.1360/N972019-00018)